



Heriot-Watt University  
Research Gateway

## Gliquidone

### Citation for published version:

Gelbrich, T, Haddow, MF & Griesser, UJ 2011, 'Gliquidone', *Acta Crystallographica Section E: Structure Reports Online*, vol. 67, no. Part 6, pp. 01343. <https://doi.org/10.1107/S1600536811016680>

### Digital Object Identifier (DOI):

[10.1107/S1600536811016680](https://doi.org/10.1107/S1600536811016680)

### Link:

[Link to publication record in Heriot-Watt Research Portal](#)

### Document Version:

Publisher's PDF, also known as Version of record

### Published In:

Acta Crystallographica Section E: Structure Reports Online

### General rights

Copyright for the publications made accessible via Heriot-Watt Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

### Take down policy

Heriot-Watt University has made every reasonable effort to ensure that the content in Heriot-Watt Research Portal complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [open.access@hw.ac.uk](mailto:open.access@hw.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

# Gliquidone

Thomas Gelbrich,\* Mairi F. Haddow† and Ulrich J. Griesser

Institute of Pharmacy, University of Innsbruck, Innrain 52, 6020 Innsbruck, Austria  
Correspondence e-mail: thomas.gelbrich@uibk.ac.at


Received 29 April 2011; accepted 3 May 2011

Key indicators: single-crystal X-ray study;  $T = 120$  K; mean  $\sigma(\text{C}—\text{C}) = 0.002$  Å;  
 $R$  factor = 0.038;  $wR$  factor = 0.101; data-to-parameter ratio = 14.2.

The title compound {systematic name: *N*-cyclohexylcarbamoyl-4-[2-(7-methoxy-4,4-dimethyl-1,3-dioxo-1,2,3,4-tetrahydroisoquinolin-2-yl)ethyl]benzenesulfonamide},  $\text{C}_{27}\text{H}_{33}\text{N}_3\text{O}_6\text{S}$ , displays an intramolecular  $\text{N}—\text{H} \cdots \text{O}=\text{S}$  interaction, as well as intermolecular  $\text{N}—\text{H} \cdots \text{O}=\text{C}$  hydrogen bonds. The latter interactions lead to the formation of hydrogen-bonded chains parallel to the  $c$  axis. The conformation of the sulfonylurea fragment is in agreement with a recent theoretical study [Kasetti *et al.* (2010). *J. Phys. Chem. B*, **114**, 11603–11610].

## Related literature

For theoretical studies of the molecular structure, see Lins *et al.* (1996); Kasetti *et al.* (2010). For thermomicroscopy, see Kuhnert-Brandstätter *et al.* (1982). For related crystal structures, see: Kobelt & Paulus (1972); Iwata *et al.* (1997); Grell *et al.* (1998); Endo *et al.* (2003).



## Experimental

### Crystal data

$\text{C}_{27}\text{H}_{33}\text{N}_3\text{O}_6\text{S}$

$M_r = 527.62$

Monoclinic,  $P2_1/c$

$a = 19.1494$  (4) Å

$b = 10.7253$  (3) Å

$c = 13.8024$  (2) Å

$\beta = 106.691$  (1)°

$V = 2715.34$  (10) Å<sup>3</sup>

$Z = 4$   
Mo  $K\alpha$  radiation  
 $\mu = 0.17$  mm<sup>-1</sup>

$T = 120$  K  
 $0.40 \times 0.40 \times 0.10$  mm

### Data collection

Bruker–Nonius Roper CCD camera  
on  $\kappa$ -goniostat diffractometer  
Absorption correction: multi-scan  
(*SADABS*; Sheldrick, 2007)  
 $T_{\min} = 0.937$ ,  $T_{\max} = 0.984$

28737 measured reflections  
5325 independent reflections  
4361 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.044$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.038$   
 $wR(F^2) = 0.101$   
 $S = 1.03$   
5325 reflections  
376 parameters  
2 restraints

H atoms treated by a mixture of  
independent and constrained  
refinement  
 $\Delta\rho_{\max} = 0.28$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.43$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å, °).

$D—H \cdots A$	$D—H$	$H \cdots A$	$D \cdots A$	$D—H \cdots A$
$\text{N1}—\text{H1N} \cdots \text{O5}^i$	0.86 (1)	2.03 (2)	2.8702 (17)	167 (2)
$\text{N2}—\text{H2N} \cdots \text{O1}$	0.87 (1)	2.15 (2)	2.8404 (17)	136 (2)

Symmetry code: (i)  $x, -y + \frac{1}{2}, z - \frac{1}{2}$ .

Data collection: *COLLECT* (Hooft, 1998); cell refinement: *DENZO* (Otwinowski & Minor, 1997) and *COLLECT*; data reduction: *DENZO* and *COLLECT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *XP* in *SHELXTL* (Sheldrick, 2008) and *Mercury* (Macrae *et al.*, 2006); software used to prepare material for publication: *publCIF* (Westrip, 2010).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: JH2285).

## References

- Endo, T., Iwata, M., Nagase, H., Shiro, M. & Ueda, H. (2003). *STP Pharma Sci.* **13**, 281–286.
- Grell, W., Hurnaus, R., Griss, G., Sauter, R., Rupprecht, E., Mark, M., Luger, P., Nar, H., Wittneben, H. & Müller, P. (1998). *J. Med. Chem.* **41**, 5219–5246.
- Hooft, R. W. W. (1998). *COLLECT*. Nonius BV, Delft, The Netherlands.
- Iwata, M., Nagase, H., Endo, T. & Ueda, H. (1997). *Acta Cryst.* **C53**, 329–331.
- Kasetti, Y., Patel, N. K., Sundriyal, S. & Bharatam, P. V. (2010). *J. Phys. Chem. B*, **114**, 11603–11610.
- Kobelt, D. & Paulus, E. F. (1972). *Acta Cryst.* **B28**, 3452–3457.
- Kuhnert-Brandstätter, M., Geiler, M. & Wurian, I. (1982). *Sci. Pharm.* **50**, 324–331.
- Lins, L., Brasseur, R. & Malaisse, W. J. (1996). *Pharm. Res.* **34**, 9–10.
- Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Sheldrick, G. M. (2007). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

† Current address: School of Chemistry, University of Bristol, Cantock's Close, Bristol BS8 1TS, England.

## supporting information

*Acta Cryst.* (2011). **E67**, o1343 [doi:10.1107/S1600536811016680]

## Gliquidone

Thomas Gelbrich, Mairi F. Haddow and Ulrich J. Griesser

### S1. Comment

Gliquidone is an anti-diabetic drug in the sulfonylurea class. It is used in the treatment of diabetes mellitus type 2. The conformation of the sulfonyl urea fragment is in agreement with the SLU-1 geometry discussed in a recent theoretical study [Kasetti *et al.* (2010)]. The same conformation was previously found in polymorph I of glimepiride (Iwata *et al.*, 1997).


In the sulfonyl urea part, the molecule displays an intramolecular N—H $\cdots$ O=S bond. The second NH group is N—H $\cdots$ O=C bonded to the imide part of a neighbouring molecule. As a result of this interaction, H-bonded chains are formed, which possess glide symmetry and propagate parallel to the *c*-axis.

### S2. Experimental

The investigated sample of gliquidone was obtained from Boehringer Ingelheim.


### S3. Refinement

All H atoms were identified in a difference map. Methyl H atoms were idealized and included as rigid groups allowed to rotate but not tip (C—H = 0.98 Å). H atoms in CH<sub>2</sub> (C—H = 0.99 Å) or CH (C—H = 1.00 Å) groups and H atoms bonded to aromatic carbon atoms (C—H = 0.95 Å) were positioned geometrically. Hydrogen atoms attached to N were refined with restrained distances [N—H = 0.88 (2) Å].  $U_{iso}$  parameters for all H atoms were refined freely.



**Figure 1**

The molecular structure with displacement ellipsoids drawn at the 50% probability level. Hydrogen atoms are shown as spheres of arbitrary size.

**Figure 2**

Fragment of an N–H $\cdots$ O=C-bonded chain. The crystal structure is viewed along [100]. H and O atoms directly engaged in H-bonds are drawn as balls.

***N*-cyclohexylcarbamoyl-4-[2-(7-methoxy-4,4-dimethyl-1,3-dioxo-1,2,3,4-tetrahydroisoquinolin-2-yl)ethyl]benzenesulfonamide**

*Crystal data*

C<sub>27</sub>H<sub>33</sub>N<sub>3</sub>O<sub>6</sub>S

$M_r = 527.62$

Monoclinic,  $P2_1/c$

Hall symbol: -P 2ybc

$a = 19.1494 (4) \text{ \AA}$

$b = 10.7253 (3) \text{ \AA}$

$c = 13.8024 (2) \text{ \AA}$

$\beta = 106.691 (1)^\circ$

$V = 2715.34 (10) \text{ \AA}^3$

$Z = 4$

$F(000) = 1120$

$D_x = 1.291 \text{ Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 16476 reflections

$\theta = 2.9\text{--}27.5^\circ$

$\mu = 0.17 \text{ mm}^{-1}$

$T = 120 \text{ K}$

Plate, colourless

$0.40 \times 0.40 \times 0.10 \text{ mm}$

*Data collection*

Bruker–Nonius Roper CCD camera on  $\kappa$ -goniostat

diffractometer

Radiation source: Bruker–Nonius FR591

rotating anode

Graphite monochromator

Detector resolution:  $9.091 \text{ pixels mm}^{-1}$

$\varphi$  and  $\omega$  scans

Absorption correction: multi-scan

(*SADABS*; Sheldrick, 2007)

$T_{\min} = 0.937$ ,  $T_{\max} = 0.984$

28737 measured reflections

5325 independent reflections

4361 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.044$

$\theta_{\max} = 26.0^\circ$ ,  $\theta_{\min} = 2.9^\circ$

$h = -23 \rightarrow 23$

$k = -13 \rightarrow 13$

$l = -15 \rightarrow 17$

*Refinement*Refinement on  $F^2$ 

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.038$  $wR(F^2) = 0.101$  $S = 1.03$ 

5325 reflections

376 parameters

2 restraints

Primary atom site location: structure-invariant  
direct methodsSecondary atom site location: difference Fourier  
mapHydrogen site location: inferred from  
neighbouring sitesH atoms treated by a mixture of independent  
and constrained refinement $w = 1/[\sigma^2(F_o^2) + (0.054P)^2 + 0.7288P]$ where  $P = (F_o^2 + 2F_c^2)/3$  $(\Delta/\sigma)_{\max} = 0.001$  $\Delta\rho_{\max} = 0.28 \text{ e } \text{\AA}^{-3}$  $\Delta\rho_{\min} = -0.43 \text{ e } \text{\AA}^{-3}$ *Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.372978 (18)	0.06096 (4)	0.34758 (2)	0.02071 (11)
O1	0.44637 (5)	0.05111 (11)	0.41217 (7)	0.0251 (2)
O2	0.35950 (6)	0.06083 (11)	0.24022 (7)	0.0291 (3)
O3	0.26601 (6)	−0.14284 (11)	0.47583 (8)	0.0301 (3)
O4	0.04953 (6)	0.56657 (10)	0.38426 (7)	0.0264 (3)
O5	0.19863 (6)	0.53754 (12)	0.70230 (8)	0.0384 (3)
O6	−0.15069 (6)	0.85561 (11)	0.38996 (8)	0.0306 (3)
N1	0.32644 (7)	−0.05733 (12)	0.37222 (9)	0.0229 (3)
H1N	0.2868 (9)	−0.0629 (18)	0.3241 (13)	0.040 (5)*
N2	0.38277 (7)	−0.07655 (13)	0.54610 (9)	0.0251 (3)
H2N	0.4223 (8)	−0.0513 (17)	0.5327 (13)	0.031 (5)*
N3	0.12334 (6)	0.55263 (11)	0.54472 (8)	0.0190 (3)
C1	0.33329 (7)	0.19457 (14)	0.38350 (10)	0.0189 (3)
C2	0.36549 (8)	0.24940 (15)	0.47723 (10)	0.0216 (3)
H2	0.4099	0.2175	0.5203	0.027 (4)*
C3	0.33191 (8)	0.35119 (15)	0.50681 (10)	0.0225 (3)
H3	0.3534	0.3886	0.5709	0.034 (5)*
C4	0.26712 (8)	0.39936 (14)	0.44388 (11)	0.0217 (3)
C5	0.23624 (8)	0.34344 (15)	0.35000 (11)	0.0244 (3)
H5	0.1922	0.3760	0.3065	0.031 (5)*
C6	0.26863 (8)	0.24152 (15)	0.31929 (11)	0.0228 (3)
H6	0.2471	0.2040	0.2553	0.031 (4)*
C7	0.22896 (8)	0.50595 (15)	0.47875 (11)	0.0248 (3)
H7A	0.2106	0.5662	0.4228	0.040 (5)*

---

H7B	0.2635	0.5500	0.5358	0.026 (4)*
C8	0.16539 (8)	0.45372 (14)	0.51243 (11)	0.0215 (3)
H8A	0.1325	0.4065	0.4558	0.026 (4)*
H8B	0.1845	0.3950	0.5692	0.031 (5)*
C9	0.06399 (7)	0.60349 (14)	0.47150 (10)	0.0184 (3)
C10	0.02037 (7)	0.69887 (13)	0.50490 (10)	0.0184 (3)
C11	0.04048 (8)	0.74227 (14)	0.60447 (10)	0.0214 (3)
C12	0.10805 (8)	0.69515 (14)	0.68274 (10)	0.0216 (3)
C13	0.14635 (8)	0.58977 (15)	0.64517 (11)	0.0232 (3)
C14	−0.04261 (7)	0.74093 (14)	0.43406 (10)	0.0204 (3)
H14	−0.0546	0.7117	0.3663	0.026 (4)*
C15	−0.08755 (8)	0.82517 (15)	0.46246 (11)	0.0235 (3)
C16	−0.06813 (9)	0.87156 (16)	0.56085 (11)	0.0301 (4)
H16	−0.0983	0.9310	0.5803	0.038 (5)*
C17	−0.00459 (9)	0.83075 (16)	0.63026 (11)	0.0290 (4)
H17	0.0086	0.8638	0.6969	0.035 (5)*
C18	0.16458 (9)	0.80078 (17)	0.71456 (13)	0.0368 (4)
H18A	0.1814	0.8253	0.6566	0.054 (6)*
H18B	0.2061	0.7716	0.7695	0.044 (5)*
H18C	0.1421	0.8727	0.7378	0.042 (5)*
C19	0.08837 (9)	0.64511 (17)	0.77617 (11)	0.0331 (4)
H19A	0.0675	0.7126	0.8069	0.047 (6)*
H19B	0.1324	0.6133	0.8254	0.039 (5)*
H19C	0.0526	0.5776	0.7556	0.068 (7)*
C20	−0.20530 (9)	0.91933 (18)	0.42316 (13)	0.0369 (4)
H20A	−0.2180	0.8696	0.4752	0.042 (5)*
H20B	−0.2488	0.9314	0.3657	0.047 (6)*
H20C	−0.1865	1.0006	0.4513	0.039 (5)*
C21	0.32254 (8)	−0.09519 (14)	0.46937 (11)	0.0217 (3)
C22	0.38491 (8)	−0.09235 (15)	0.65249 (11)	0.0228 (3)
H22	0.3337	−0.1030	0.6557	0.031 (4)*
C23	0.42853 (9)	−0.20722 (15)	0.69978 (12)	0.0292 (4)
H23A	0.4786	−0.2015	0.6932	0.029 (4)*
H23B	0.4053	−0.2830	0.6636	0.049 (6)*
C24	0.43184 (9)	−0.21684 (16)	0.81155 (12)	0.0333 (4)
H24A	0.3821	−0.2302	0.8174	0.041 (5)*
H24B	0.4619	−0.2897	0.8419	0.051 (6)*
C25	0.46396 (9)	−0.10010 (17)	0.86938 (12)	0.0328 (4)
H25A	0.5152	−0.0906	0.8687	0.032 (5)*
H25B	0.4636	−0.1082	0.9407	0.039 (5)*
C26	0.42077 (9)	0.01468 (17)	0.82278 (12)	0.0314 (4)
H26A	0.4446	0.0901	0.8588	0.035 (5)*
H26B	0.3710	0.0096	0.8304	0.039 (5)*
C27	0.41609 (8)	0.02515 (15)	0.71081 (11)	0.0251 (3)
H27A	0.3849	0.0972	0.6812	0.031 (5)*
H27B	0.4654	0.0407	0.7039	0.027 (4)*

---

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
S1	0.01746 (19)	0.0296 (2)	0.01411 (19)	0.00384 (15)	0.00301 (14)	−0.00118 (14)
O1	0.0158 (5)	0.0382 (7)	0.0206 (5)	0.0056 (4)	0.0038 (4)	0.0007 (4)
O2	0.0295 (6)	0.0427 (7)	0.0148 (5)	0.0065 (5)	0.0059 (4)	−0.0014 (5)
O3	0.0242 (6)	0.0330 (7)	0.0310 (6)	−0.0057 (5)	0.0048 (4)	−0.0019 (5)
O4	0.0293 (6)	0.0314 (6)	0.0161 (5)	0.0073 (5)	0.0026 (4)	−0.0029 (4)
O5	0.0346 (7)	0.0451 (8)	0.0252 (6)	0.0187 (6)	−0.0080 (5)	−0.0040 (5)
O6	0.0235 (6)	0.0404 (7)	0.0238 (5)	0.0148 (5)	0.0003 (4)	0.0004 (5)
N1	0.0215 (7)	0.0252 (7)	0.0183 (6)	0.0010 (5)	0.0000 (5)	−0.0041 (5)
N2	0.0173 (6)	0.0384 (8)	0.0185 (6)	0.0005 (6)	0.0033 (5)	0.0028 (5)
N3	0.0191 (6)	0.0196 (6)	0.0169 (6)	0.0035 (5)	0.0029 (5)	0.0004 (5)
C1	0.0166 (7)	0.0237 (8)	0.0166 (7)	−0.0004 (6)	0.0054 (5)	0.0004 (6)
C2	0.0157 (7)	0.0307 (8)	0.0164 (7)	−0.0004 (6)	0.0014 (5)	0.0014 (6)
C3	0.0219 (7)	0.0284 (8)	0.0166 (7)	−0.0022 (6)	0.0045 (6)	−0.0036 (6)
C4	0.0207 (7)	0.0228 (8)	0.0229 (7)	−0.0010 (6)	0.0085 (6)	0.0005 (6)
C5	0.0190 (7)	0.0285 (9)	0.0219 (7)	0.0036 (6)	−0.0002 (6)	0.0010 (6)
C6	0.0201 (7)	0.0268 (8)	0.0182 (7)	0.0006 (6)	0.0002 (6)	−0.0019 (6)
C7	0.0242 (8)	0.0250 (8)	0.0257 (8)	0.0009 (6)	0.0078 (6)	−0.0015 (6)
C8	0.0211 (7)	0.0204 (8)	0.0216 (7)	0.0043 (6)	0.0042 (6)	0.0003 (6)
C9	0.0183 (7)	0.0198 (7)	0.0161 (7)	−0.0006 (6)	0.0032 (5)	0.0016 (6)
C10	0.0193 (7)	0.0182 (7)	0.0166 (7)	−0.0001 (6)	0.0034 (5)	0.0012 (5)
C11	0.0238 (8)	0.0219 (8)	0.0166 (7)	0.0007 (6)	0.0024 (6)	0.0002 (6)
C12	0.0242 (7)	0.0216 (8)	0.0161 (7)	0.0021 (6)	0.0009 (6)	−0.0008 (6)
C13	0.0206 (7)	0.0270 (8)	0.0185 (7)	0.0018 (6)	−0.0001 (6)	0.0003 (6)
C14	0.0208 (7)	0.0237 (8)	0.0153 (7)	0.0013 (6)	0.0031 (5)	0.0008 (6)
C15	0.0211 (7)	0.0254 (8)	0.0212 (7)	0.0060 (6)	0.0015 (6)	0.0032 (6)
C16	0.0332 (9)	0.0292 (9)	0.0261 (8)	0.0125 (7)	0.0056 (7)	−0.0037 (7)
C17	0.0347 (9)	0.0293 (9)	0.0193 (7)	0.0091 (7)	0.0018 (6)	−0.0046 (6)
C18	0.0340 (9)	0.0289 (9)	0.0375 (9)	−0.0039 (8)	−0.0057 (7)	−0.0019 (7)
C19	0.0365 (9)	0.0412 (10)	0.0205 (8)	0.0113 (8)	0.0064 (7)	0.0078 (7)
C20	0.0274 (9)	0.0439 (11)	0.0387 (10)	0.0177 (8)	0.0084 (7)	0.0050 (8)
C21	0.0221 (8)	0.0189 (7)	0.0229 (7)	0.0044 (6)	0.0045 (6)	−0.0022 (6)
C22	0.0169 (7)	0.0304 (9)	0.0207 (7)	0.0018 (6)	0.0048 (6)	0.0058 (6)
C23	0.0310 (9)	0.0233 (8)	0.0313 (8)	0.0007 (7)	0.0059 (7)	0.0023 (7)
C24	0.0335 (9)	0.0319 (9)	0.0333 (9)	0.0044 (7)	0.0074 (7)	0.0139 (7)
C25	0.0330 (9)	0.0425 (10)	0.0210 (8)	0.0055 (8)	0.0044 (6)	0.0052 (7)
C26	0.0348 (9)	0.0357 (10)	0.0246 (8)	0.0082 (8)	0.0101 (7)	0.0001 (7)
C27	0.0264 (8)	0.0242 (8)	0.0251 (8)	0.0061 (6)	0.0078 (6)	0.0040 (6)

*Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )*

S1—O2	1.4293 (10)	C11—C12	1.5146 (19)
S1—O1	1.4357 (10)	C12—C13	1.517 (2)
S1—N1	1.6413 (14)	C12—C19	1.540 (2)
S1—C1	1.7588 (15)	C12—C18	1.541 (2)
O3—C21	1.2234 (18)	C14—C15	1.380 (2)

O4—C9	1.2213 (17)	C14—H14	0.9500
O5—C13	1.2184 (18)	C15—C16	1.393 (2)
O6—C15	1.3692 (17)	C16—C17	1.386 (2)
O6—C20	1.4305 (19)	C16—H16	0.9500
N1—C21	1.4233 (19)	C17—H17	0.9500
N1—H1N	0.856 (14)	C18—H18A	0.9800
N2—C21	1.3380 (19)	C18—H18B	0.9800
N2—C22	1.4670 (18)	C18—H18C	0.9800
N2—H2N	0.872 (14)	C19—H19A	0.9800
N3—C13	1.3871 (18)	C19—H19B	0.9800
N3—C9	1.3968 (18)	C19—H19C	0.9800
N3—C8	1.4759 (18)	C20—H20A	0.9800
C1—C2	1.393 (2)	C20—H20B	0.9800
C1—C6	1.3935 (19)	C20—H20C	0.9800
C2—C3	1.387 (2)	C22—C27	1.522 (2)
C2—H2	0.9500	C22—C23	1.526 (2)
C3—C4	1.393 (2)	C22—H22	1.0000
C3—H3	0.9500	C23—C24	1.529 (2)
C4—C5	1.396 (2)	C23—H23A	0.9900
C4—C7	1.508 (2)	C23—H23B	0.9900
C5—C6	1.382 (2)	C24—C25	1.517 (2)
C5—H5	0.9500	C24—H24A	0.9900
C6—H6	0.9500	C24—H24B	0.9900
C7—C8	1.529 (2)	C25—C26	1.519 (2)
C7—H7A	0.9900	C25—H25A	0.9900
C7—H7B	0.9900	C25—H25B	0.9900
C8—H8A	0.9900	C26—C27	1.526 (2)
C8—H8B	0.9900	C26—H26A	0.9900
C9—C10	1.476 (2)	C26—H26B	0.9900
C10—C14	1.3916 (19)	C27—H27A	0.9900
C10—C11	1.3963 (19)	C27—H27B	0.9900
C11—C17	1.396 (2)		
O2—S1—O1	119.79 (6)	O6—C15—C14	116.12 (13)
O2—S1—N1	105.48 (7)	O6—C15—C16	124.05 (14)
O1—S1—N1	107.90 (7)	C14—C15—C16	119.82 (13)
O2—S1—C1	109.21 (7)	C17—C16—C15	119.79 (14)
O1—S1—C1	108.07 (7)	C17—C16—H16	120.1
N1—S1—C1	105.49 (7)	C15—C16—H16	120.1
C15—O6—C20	116.95 (12)	C16—C17—C11	121.52 (14)
C21—N1—S1	126.46 (10)	C16—C17—H17	119.2
C21—N1—H1N	115.9 (13)	C11—C17—H17	119.2
S1—N1—H1N	107.8 (13)	C12—C18—H18A	109.5
C21—N2—C22	123.02 (13)	C12—C18—H18B	109.5
C21—N2—H2N	118.8 (11)	H18A—C18—H18B	109.5
C22—N2—H2N	118.1 (11)	C12—C18—H18C	109.5
C13—N3—C9	124.75 (12)	H18A—C18—H18C	109.5
C13—N3—C8	117.64 (11)	H18B—C18—H18C	109.5



C9—N3—C8	117.60 (11)	C12—C19—H19A	109.5
C2—C1—C6	120.89 (14)	C12—C19—H19B	109.5
C2—C1—S1	119.59 (11)	H19A—C19—H19B	109.5
C6—C1—S1	119.48 (11)	C12—C19—H19C	109.5
C3—C2—C1	119.12 (13)	H19A—C19—H19C	109.5
C3—C2—H2	120.4	H19B—C19—H19C	109.5
C1—C2—H2	120.4	O6—C20—H20A	109.5
C2—C3—C4	120.90 (13)	O6—C20—H20B	109.5
C2—C3—H3	119.6	H20A—C20—H20B	109.5
C4—C3—H3	119.6	O6—C20—H20C	109.5
C3—C4—C5	118.90 (14)	H20A—C20—H20C	109.5
C3—C4—C7	120.55 (13)	H20B—C20—H20C	109.5
C5—C4—C7	120.47 (13)	O3—C21—N2	125.76 (14)
C6—C5—C4	121.11 (13)	O3—C21—N1	118.35 (13)
C6—C5—H5	119.4	N2—C21—N1	115.88 (13)
C4—C5—H5	119.4	N2—C22—C27	109.13 (12)
C5—C6—C1	119.07 (13)	N2—C22—C23	112.08 (12)
C5—C6—H6	120.5	C27—C22—C23	111.03 (12)
C1—C6—H6	120.5	N2—C22—H22	108.2
C4—C7—C8	108.68 (12)	C27—C22—H22	108.2
C4—C7—H7A	110.0	C23—C22—H22	108.2
C8—C7—H7A	110.0	C22—C23—C24	110.18 (13)
C4—C7—H7B	110.0	C22—C23—H23A	109.6
C8—C7—H7B	110.0	C24—C23—H23A	109.6
H7A—C7—H7B	108.3	C22—C23—H23B	109.6
N3—C8—C7	112.30 (12)	C24—C23—H23B	109.6
N3—C8—H8A	109.1	H23A—C23—H23B	108.1
C7—C8—H8A	109.1	C25—C24—C23	111.60 (13)
N3—C8—H8B	109.1	C25—C24—H24A	109.3
C7—C8—H8B	109.1	C23—C24—H24A	109.3
H8A—C8—H8B	107.9	C25—C24—H24B	109.3
O4—C9—N3	119.59 (13)	C23—C24—H24B	109.3
O4—C9—C10	122.97 (12)	H24A—C24—H24B	108.0
N3—C9—C10	117.42 (12)	C24—C25—C26	110.91 (13)
C14—C10—C11	121.46 (13)	C24—C25—H25A	109.5
C14—C10—C9	117.34 (12)	C26—C25—H25A	109.5
C11—C10—C9	121.18 (12)	C24—C25—H25B	109.5
C17—C11—C10	117.50 (13)	C26—C25—H25B	109.5
C17—C11—C12	120.60 (12)	H25A—C25—H25B	108.0
C10—C11—C12	121.90 (13)	C25—C26—C27	110.78 (13)
C11—C12—C13	113.74 (11)	C25—C26—H26A	109.5
C11—C12—C19	110.43 (12)	C27—C26—H26A	109.5
C13—C12—C19	106.57 (12)	C25—C26—H26B	109.5
C11—C12—C18	110.13 (13)	C27—C26—H26B	109.5
C13—C12—C18	106.03 (13)	H26A—C26—H26B	108.1
C19—C12—C18	109.78 (13)	C22—C27—C26	112.00 (13)
O5—C13—N3	118.75 (14)	C22—C27—H27A	109.2
O5—C13—C12	120.60 (13)	C26—C27—H27A	109.2

N3—C13—C12	120.63 (12)	C22—C27—H27B	109.2
C15—C14—C10	119.86 (13)	C26—C27—H27B	109.2
C15—C14—H14	120.1	H27A—C27—H27B	107.9
C10—C14—H14	120.1		

*Hydrogen-bond geometry (Å, °)*

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
N1—H1N $\cdots$ O5 <sup>i</sup>	0.86 (1)	2.03 (2)	2.8702 (17)	167 (2)
N2—H2N $\cdots$ O1	0.87 (1)	2.15 (2)	2.8404 (17)	136 (2)

Symmetry code: (i)  $x, -y+1/2, z-1/2$ .